

Expression and Characterization of HIV-1 Envelope Protein Associated with a
Broadly Reactive Neutralizing Antibody Response

5 Inventors: Gerald V. Quinnan, Jr. & Peng Fei Zhang

Related Applications

This application is related to U.S. Provisional Application 60/095,267 filed
10 August 4, 1998, which is herein incorporated by reference in its entirety.

Technical Field

The present invention relates to HIV-1 envelope proteins and peptides derived
15 from the donor of the Neutralizing Reference Human Serum (2) which is noted for its
capacity to neutralize primary HIV isolates of varied subtypes.

Acknowledgment of Federal Support

20 The present invention arose in part from research funded by the following federal
grant monies: NIH AI37436 and AI44339, and USUHS R087E2

Background of the Invention

25 The development of a successful vaccine against HIV infection or a vaccine agent
capable of preventing HIV disease progression has been a public health goal for over 15
years. One of the immune responses that may be required to elicit a protective immune
response against HIV infection is the generation of antibodies that are virus neutralizing.

The target of HIV-1 neutralizing antibodies (NA) is the envelope glycoprotein
30 complex. This complex is a multimeric structure composed of three or four copies each
of the gp120 surface and gp41 transmembrane glycoproteins (Luciw, 1996). There are a

number of neutralization domains on each of the three or four heterodimeric components of the complex (Thali *et al.*, 1992, 1993; Zwart *et al.*, 1991; Moore *et al.*, 1993; Trkola *et al.*, 1996; Muster *et al.*, 1993; Cotropia *et al.*, 1996; Sabri *et al.*, 1996). The amino acid compositions of the proteins vary substantially from strain to strain. Some of the

5 neutralization domains are in regions which tend to vary greatly, while others are in regions which tend to be highly conserved. The variable neutralization domains include those in variable (V) regions 1, 2, and 3 of gp120, while the conserved domains include the primary receptor binding site, and other epitopes in gp120 and gp41. Amino acid sequence variation is undoubtedly the explanation for the variation that is seen in

10 specificity of neutralization sensitivity among virus strains. However, it has not been possible to classify antigenic subtypes of HIV-1 based on genetic analyses, and various regions of the envelope complex even outside of the neutralization domains have been shown to contribute to antigenic variability (Thali *et al.*, 1994; Back *et al.*, 1993).

Recent findings indicate that the neutralization of primary isolates of HIV may be

15 mediated primarily by antibodies directed against non-V3 region epitopes expressed on the oligomeric complex but not on monomeric gp120, while laboratory adapted strains are more readily neutralized by antibodies directed against V3 (Hioe *et al.*, 1997; VanCott *et al.*, 1997). The identity of the non-V3 epitopes recognized on primary isolates is not established. The presence of antibodies which have broadly neutralizing activity against

20 primary isolates of many subtypes of HIV-1 in sera from infected people is unusual, but the nature of the envelope proteins in individuals with such antibodies may be of interest for defining the epitopes which may be broadly immunogenic in vaccines.

25 Summary of the Invention

The present inventors have cloned and characterized the envelope genes from the donor of human serum which is noted for its capacity to neutralize primary HIV isolates of various subtype (Vujcic, *et al.* 1995, D'Souza *et al.*, 1991).

The invention includes an isolated HIV envelope protein or fragment thereof

30 which, when injected into a mammal, induces the production of broadly cross-reactive neutralizing anti-serum against multiple strains of HIV-1.

The invention further includes an isolated HIV envelope protein or fragment thereof comprising a proline at a position corresponding to amino acid residue 313, a methionine at a position corresponding to amino acid residue 314 and a glutamine at a position corresponding to amino acid residue 325 of SEQ ID NO:1.

5 In another embodiment, the invention includes an isolated HIV envelope glycoprotein or fragment thereof comprising an alanine at a position corresponding to amino acid residue 167 of SEQ ID NO:1.

The invention also includes an isolated HIV envelope protein comprising the amino acid sequence of SEQ ID NO:1 as well as an isolated nucleic acid molecule
10 encoding the envelope protein.

Compositions for eliciting an immune response, such as vaccines, immunogenic compositions and attenuated viral vaccine delivery vectors comprising the envelope proteins, peptides and nucleic acids encoding such proteins and peptides of the invention are also included. Methods for generating antibodies in a mammal comprising
15 administering one or more of these proteins, peptides and nucleic acids, in an amount sufficient to induce the production of the antibodies, is also included in the invention.

The invention also comprises a diagnostic reagent comprising one or more of the isolated HIV-1 envelope proteins and methods for detecting broadly cross-reactive neutralizing anti-serum against multiple strains of HIV-1.

20

Brief Description of the Drawings

Figure 1: Phylogenetic analysis of the gp120 and gp41 nucleotide coding sequences of clone R2. Alignments were performed using the Clustal algorithm of
25 Higgins and Sharp in the program DNA Star (Higgins *et al.*, 1989; Saitou *et al.*, 1987; Myers *et al.*, 1988). The graphs at the bottom of the two figures indicate the percent similarity distances represented by the dendograms. Gene bank accession numbers for the sequences represented are: MW 959, U08453; MW960, U08454; D747, X65638; BR020, U27401; BR029, U27413; RU131, U30312; UG975, U27426; AD8, M60472;
30 HXB, K03455; NDK, M27323; Z2Z6, M22639; UG021, U27399; CM235, L03698;

TH022, U09139; TH006, U08810; UG275, L22951; SF1703, M66533; RW020, U08794; RW009, U08793; U455, M62320; and Z321, M15896.

Figure 2: Neutralization of clade B viruses and pseudoviruses by sera from 10 male residents of the Baltimore/Washington, D.C. area collected from 1985-1990 in the Multicenter AIDS Cohort Study. The P9 and P10 viruses (P9-V and P10-V) are primary isolates from two of the serum donors (Quinnan *et al.*, 1998). The neutralization assays were performed in PM1 cells, as described in the Examples. Each point represents the results obtained with an individual serum. The open bars represent the standard deviations about the geometric means, indicated by the midlines. The numbers above the results obtained using pseudoviruses indicate the probabilities obtained from testing the null hypothesis by paired t testing comparing the individual pseudoviruses to R2.

Subject
~~Figure 3 (A): Inhibition of Reference 2-mediated neutralization of pseudoviruses by synthetic V3 peptides. The neutralization endpoints for 90% neutralization were calculated as described previously (Quinnan *et al.*, 1999; Quinnan *et al.*, 1998; Zhang *et al.*, 1999; Park *et al.*, 1998). Results shown are means of triplicate determinations. Dose-response effects of R2 linear 17-mer (open square) and cyclic (closed square) and the 93TH966.8 cyclic (shaded square) V3 peptides on neutralization of clone R2 pseudovirus. The peptide concentrations are 3×10 raised to the indicated power.~~

Figure 3 (B): Comparative inhibitory effects of peptides on neutralization of R2 and MN (clone V5) pseudoviruses. All peptides were tested at 15 $\mu\text{g/ml}$. The linear peptides (L) corresponded to the apical sequences of the respective V3 loops. The cyclic peptides (C) corresponded to the full lengths of the respective V3 regions of the different strains. Neutralization in the absence of peptide (None), is also shown.

Figure 4 (A): Effect of cyclic R2 V3 peptide on neutralization of pseudoviruses. Fold inhibition of neutralization was calculated as the ratio of the 50% neutralization titer obtained in the absence of peptide compared to that obtained in the presence of cyclic R2 V3 peptide (15 $\mu\text{g/ml}$). All assays were performed in triplicate. Neutralization titers

were calculated at the midpoints of the infectivity inhibition curves, since the curves tended to be most parallel in this region. Similar results were obtained comparing 90% neutralization endpoints. Peptide inhibition of neutralization of R2 pseudovirus by sera from MACS donors (donor numbers 1-10), two assays each, and by Reference 2. Results are shown for two determinations for each serum from the MACS donors and for 12 assays of Reference 2 performed during the same time intervals as the other experiments shown in panels (A) and (B).

Figure 4 (B): Peptide inhibition of neutralization of pseudoviruses expressing MACS patient envelopes (patient numbers 3, 4, 6, 8, 9, and 10) by Reference 2. Results of two or three separate assays of each pseudovirus are shown.

Modes of Carrying Out the Invention

General Description

A goal of immunization against HIV is to induce neutralizing antibody (NA) responses broadly reactive against diverse strains of virus. The present inventors have studied envelope protein from a donor with non-progressive HIV-1 infection whose serum contains broadly cross-reactive, primary virus NA. DNA was extracted from lymphocytes, which had been collected approximately six and twelve months prior to the time of collection of the cross reactive serum, *env* genes were synthesized by nested PCR, cloned, expressed on pseudoviruses, and phenotyped in NA assays. Two clones from each time point had identical V3 region nucleotide sequences, utilized CCR5 but not CXCR4 for cell entry, and had similar reactivities with two reference sera. Analysis of the full nucleotide sequence of one clone demonstrated it to be subtype B, with a predicted GPGRFV apical V3 sequence, normal predicted glycosylation, and an intact reading frame. Infectivity assays of R2 pseudovirus in HOS cells expressing CD4 and various coreceptors demonstrated the envelope to be CCR5 dependent. R2 pseudovirus was compared to others expressing *env* genes of various clades for neutralization by sera from donors in the United States (presumed or known subtype B infections), and from individuals infected with subtypes A, C, and E viruses. Neutralization by the sera from

donors in the United States of pseudoviruses expressing R2 and other clade B envs was similarly low to moderate, although R2 was uniquely neutralized by all. R2 was neutralized by sera from people infected with clades A-F, while other clade B, D, E and G pseudoviruses were neutralized less often. One highly sensitive clade C pseudovirus was neutralized by all the sera, although the titers varied more than 250-fold. The results suggest that the epitope(s) which induced the cross-clade reactive NA in Donor 2 may be expressed on the R2 envelope.

The present invention relates to HIV-1 envelope proteins from this donor who had non-progressive HIV-1 infection whose serum contains broadly cross-reactive, primary virus neutralizing antibody. The invention also relates to isolated or purified proteins and protein fragments that share certain amino acids at particular positions with the foregoing HIV-1 proteins.

15 Specific Embodiments

Proteins and Peptides

Proteins and peptides of the invention include the full length envelope protein having the amino acid sequence of Table 3 (SEQ ID NO:1), gp120 having the amino acid sequence corresponding to gp120 in Table 3 (amino acids: 1-520 of SEQ ID NO:1), gp41 having the amino acid sequence corresponding to gp41 in Table 3 (amino acids 521-866 of SEQ ID NO:1), as well as polypeptides and peptides corresponding to the V3 domain and other domains such as V1/V2, C3, V4, C4 and V5. These domains correspond to the following amino acid residues of SEQ ID NO:1:

25

DOMAIN	AMINO ACID RESIDUES
C1	30-124
V1	125-162
V2	163-201
C2	202-300
V3	301-336
C3	337-387

30

DOMAIN	AMINO ACID RESIDUES
V4	388-424
C4	425-465
V5	466-509
C5	510-520

5

Sub C2

SEQUENCE LISTING

- ~~Polypeptides and peptides comprising any single domain may be of variable length but include the amino acid residues of Table 3 (SEQ ID NO:1) which differ from previously sequenced envelope proteins. For instance, peptides of the invention which include all or part of the V3 domain may comprise the sequence: PM X₁ X₂ X₃ X₄ X₅ X₆ X₇ X₈ X₉ X₁₀ Q, wherein X₁-X₁₀ are any natural or non-natural amino acids (P refers to Proline, M refers to methionine and Q refers to Glutamine). Non-natural amino acids include, for example, beta-alanine (beta-Ala), or other omega-amino acids, such as 3-amino propionic, 2,3-diamino propionic (2,3-diaP), 4-amino butyric and so forth, alpha-aminisobutyric acid (Aib), sarcosine (Sat), ornithine (Orn), citrulline (Cit), t-butylalanine (t-BuA), t-butylglycine (t-BuG), N-methylisoleucine (N-MeIle), phenylglycine (Phg), and cyclohexylalanine (Cha), norleucine (Nle), cysteic acid (Cya) 2-naphthylalanine (2-Nal); 1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid (Tic); beta-2-thienylalanine (Thi); and methionine sulfoxide (MSO). Preferably, peptides of the invention are 60%, 70%, 80% or more preferably, 90% identical to the V3 region of the HIV envelope protein of Table 3 (SEQ ID NO:1). Accordingly, V3 peptides of the invention comprise about 13 amino acids but may be 14, 15, 17, 20, 25, 30, 35, 36, 39, 40, 45, 50 or more amino acids in length. In one embodiment, a V3 peptide of 13 amino acids in length consists of the sequence PMGPGRAFYTGTGQ (amino acids 313-325 of Table 3 (SEQ ID NO:1)).~~

- 25 In another embodiment of the invention, polypeptides and peptides comprising all or part of the V1/V2 domain comprise an amino acid sequence with an alanine residue at a position corresponding to amino acid 167 Table 3 (SEQ ID NO:1). For instance, peptides of the invention spanning the V1/V2 domain may comprise the sequence ENIATSIG (residues 164-171 of SEQ ID NO:1) and may be about 8, 9, 10, 15, 20, 25,

30, 35, 40, 45, 50 or more amino acids in length. As used herein, "at a position corresponding to" refers to amino acid positions in HIV envelope proteins or peptides of the invention which are equivalent to a given amino acid residue in the sequence of Table 1 (SEQ ID NO:1) in the context of the surrounding residues.

5 The peptides of the present invention may be prepared by any known techniques. Conveniently, the peptides may be prepared using the solid-phase synthetic technique initially described by Merrifield (1965), which is incorporated herein by reference. Other peptide synthesis techniques may be found, for example, in Bodanszky *et al.*, *Peptide Synthesis*, 2d ed. (New York, Wiley, 1976).

10

Nucleic acids and Recombinant Expression of Peptide or Proteins

Proteins and peptides of the invention may be prepared by any available means, including recombinant expression of the desired protein or peptide in eukaryotic or prokaryotic host cells (see U.S. Patent 5,696,238). Methods for producing proteins or peptides of the invention for purification may employ conventional molecular biology, microbiology, and recombinant DNA techniques within the ordinary skill level of the art. Such techniques are explained fully in the literature. See, for example, Maniatis *et al.*, *Molecular Cloning: A Laboratory Manual*, 2d ed. (Cold Spring Harbor, Cold Spring Harbor Laboratory Press, 1989); Glover, *DNA Cloning: A Practical Approach*, Vols. 1-4 (Oxford, IRL Press, 1985); Gait, *Oligonucleotide Synthesis: A Practical Approach* (Oxford, IRL Press, 1984); Hames & Higgins, *Nucleic Acid Hybridisation: A Practical Approach* (Oxford, IRL Press, 1985); Freshney, *Animal Cell Culture: A Practical Approach* (Oxford, IRL Press, 1992); Perbal, *A Practical Guide To Molecular Cloning* (New York, Wiley, 1984).

25 The present invention further provides nucleic acid molecules that encode the proteins or peptides of the invention. Such nucleic acid molecules can be in an isolated form, or can be operably linked to expression control elements or vector sequences. The present invention further provides host cells that contain the vectors via transformation, transfection, electroporation or any other art recognized means of introducing a nucleic acid into a cell.

30

As used herein, a "cell line" is a clone of a primary cell that is capable of stable growth *in vitro* for many generations.

A DNA "coding sequence" is a double-stranded DNA sequence which is transcribed and translated into a polypeptide *in vivo* when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxy) terminus. A polyadenylation signal and transcription termination sequence will usually be located 3' to the coding sequence.

A "heterologous" region of the DNA construct is an identifiable segment of DNA within a larger DNA molecule that is not found in association with the larger molecule in nature. Thus, when the heterologous region encodes a mammalian gene, the gene will usually be flanked by DNA that does not flank the mammalian genomic DNA in the genome of the source organism. Another example of a heterologous coding sequence is a construct where the coding sequence itself is not found in nature (*e.g.*, a cDNA where the genomic coding sequence contains introns, or synthetic sequences having codons different than the native gene). Allelic variations or naturally-occurring mutational events do not give rise to a heterologous region of DNA as defined herein.

As used herein, "naked DNA" means nucleic acid molecules that are free from viral particles, particularly retroviral particles. This term also means nucleic acid molecules which are free from facilitator agents including but not limited to the group comprising: lipids, liposomes, extracellular matrix-active enzymes, saponins, lectins, estrogenic compounds and steroidal hormones, hydroxylated lower alkyls, dimethyl sulfoxide (DMSO) and urea.

As used herein, a "nucleic acid molecule" refers to the polymeric form of deoxyribonucleotides (adenine, guanine, thymine, and/or cytosine) in either its single stranded form, or in double-stranded helix as well as RNA. This term refers only to the primary and secondary structure of the molecule and is not limited to any particular tertiary form. In discussing the structure of particular double-stranded DNA molecules, sequences may be described herein according to the normal convention of giving only the sequence in the 5' to 3' direction along the nontranscribed strand of DNA (*e.g.*, the strand having a sequence homologous to the mRNA). Transcriptional and translational control

sequences are DNA regulatory sequences, such as promoters, enhancers, polyadenylation signals, terminators, and the like, that provide for the expression of a coding sequence in a host-cell.

As used herein, a "promoter sequence" is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. For purposes of defining the present invention, the promoter sequence is bounded (inclusively) at its 3' terminus by the transcription initiation site and extends upstream (5' direction) to include the minimum number of bases or elements necessary to initiate transcription at levels detectable above background. Within the promoter sequence will be found a transcription initiation site, as well as protein binding domains responsible for the binding of RNA polymerase. Eukaryotic promoters will often, but not always, contain "TATA" boxes and "CAT" boxes.

As used herein, a "replicon" is any genetic element (*e.g.*, plasmid, chromosome, virus) that functions as an autonomous unit of DNA replication *in vivo*; *i.e.*, capable of replication under its own control.

A "signal sequence" can be included before the coding sequence or the native 29 amino acid signal sequence from the envelope protein of Table 3 may be used. This sequence encodes a signal peptide, N-terminal to the polypeptide, that communicates to the host cell to direct the polypeptide to the cell surface or secrete the polypeptide into the media. This signal peptide is clipped off by the host cell before the protein leaves the cell. Signal sequences can be found associated with a variety of proteins native to prokaryotes and eukaryotes. For instance, alpha-factor, a native yeast protein, is secreted from yeast, and its signal sequence can be attached to heterologous proteins to be secreted into the media (See U.S. Pat. No. 4,546,082, and EP 0116201). Further, the alpha-factor and its analogs have been found to secrete heterologous proteins from a variety of yeast, such as *Saccharomyces* and *Kluyveromyces*, (EP 88312306.9; EP 0324274 publication, and EP 0301669). An example for use in mammalian cells is the tPA signal used for expressing Factor VIIIc light chain.

As used herein, DNA sequences are "substantially homologous" when at least about 85% (preferably at least about 90% and most preferably at least about 95%) of the nucleotides match over the defined length of the DNA sequences. Sequences that are

substantially homologous can be identified in a Southern hybridization experiment under, for example, stringent conditions as defined for that particular system. Defining appropriate hybridization conditions is within the skill of the art. See, for example, Maniatis *et al.*, *supra*.

- 5 A cell has been "transformed" by exogenous or heterologous DNA when such DNA has been introduced inside the cell. The transforming DNA may or may not be integrated (covalently linked) into chromosomal DNA making up the genome of the cell. In prokaryotes, for example, the transforming DNA may be maintained on an episomal element such as a plasmid or viral vector. With respect to eukaryotic cells, a stably
- 10 transformed cell is one in which the transforming DNA has become integrated into a chromosome so that it is inherited by daughter cells through chromosome replication. This stability is demonstrated by the ability of the eukaryotic cell to establish cell lines or clones comprised of a population of daughter cells containing the transforming DNA.

- A coding sequence is "under the control" of transcriptional and translational
- 15 control sequences in a cell when RNA polymerase transcribes the coding sequence into mRNA, which is then translated into the protein encoded by the coding sequence.

 As used herein, a "vector" is a replicon, such as plasmid, phage or cosmid, to which another DNA segment may be attached so as to bring about the replication of the attached segment.

- 20 Vectors are used to simplify manipulation of the DNA which encodes the HIV proteins or peptides, either for preparation of large quantities of DNA for further processing (cloning vectors) or for expression of the HIV proteins or peptides (expression vectors). Vectors comprise plasmids, viruses (including phage), and integrated DNA fragments, *i.e.*, fragments that are integrated into the host genome by recombination.
- 25 Cloning vectors need not contain expression control sequences. However, control sequences in an expression vector include transcriptional and translational control sequences such as a transcriptional promoter, a sequence encoding suitable ribosome binding sites, and sequences which control termination of transcription and translation. The expression vector should preferably include a selection gene to facilitate the stable
- 30 expression of HIV gene and/or to identify transformants. However, the selection gene for

maintaining expression can be supplied by a separate vector in cotransformation systems using eukaryotic host cells.

Suitable vectors generally will contain replicon (origins of replication, for use in non-integrative vectors) and control sequences which are derived from species compatible with the intended expression host. By the term "replicable" vector as used herein, it is intended to encompass vectors containing such replicons as well as vectors which are replicated by integration into the host genome. Transformed host cells are cells which have been transformed or transfected with vectors containing HIV peptide or protein encoding DNA. The expressed HIV proteins or peptides may be secreted into the culture supernatant, under the control of suitable processing signals in the expressed peptide, *e.g.* homologous or heterologous signal sequences.

Expression vectors for host cells ordinarily include an origin of replication, a promoter located upstream from the HIV protein or peptide coding sequence, together with a ribosome binding site, a polyadenylation site, and a transcriptional termination sequence. Those of ordinary skill will appreciate that certain of these sequences are not required for expression in certain hosts. An expression vector for use with microbes need only contain an origin of replication recognized by the host, a promoter which will function in the host, and a selection gene.

Commonly used promoters are derived from polyoma, bovine papilloma virus, CMV (cytomegalovirus, either murine or human), Rouse sarcoma virus, adenovirus, and simian virus 40 (SV40). Other control sequences (*e.g.*, terminator, polyA, enhancer, or amplification sequences) can also be used.

An expression vector is constructed so that the HIV protein or peptide coding sequence is located in the vector with the appropriate regulatory sequences, the positioning and orientation of the coding sequence with respect to the control sequences being such that the coding sequence is transcribed and translated under the "control" of the control sequences (*i.e.*, RNA polymerase which binds to the DNA molecule at the control sequences transcribes the coding sequence). The control sequences may be ligated to the coding sequence prior to insertion into a vector, such as the cloning vectors described above. Alternatively, the coding sequence can be cloned directly into an expression vector which already contains the control sequences and an appropriate

restriction site. If the selected host cell is a mammalian cell, the control sequences can be heterologous or homologous to the HIV coding sequence, and the coding sequence can either be genomic DNA containing introns or cDNA.

Higher eukaryotic cell cultures may be used to express the proteins of the present invention, whether from vertebrate or invertebrate cells, including insects, and the procedures of propagation thereof are known. See, for example, Kruse & Patterson, *Tissue Culture* (New York, Academic Press, 1973).

Suitable host cells for expressing HIV proteins or peptides in higher eukaryotes include: monkey kidney CVI line transformed by SV40 (COS-7, ATCC CRL1651); baby hamster kidney cells (BHK, ATCC CRL10); Chinese hamster ovary cells-DHFR (Urlaub & Chasin, 1980); mouse Sertoli cells (Mather, 1980); monkey kidney cells (CVI ATCC CCL70); African green monkey kidney cells (VERO76, ATCC CRL1587); human cervical carcinoma cells (HeLa, ATCC CCL2); canine kidney cells (MDCK, ATCC CCL34); buffalo rat liver cells (BRL3A, ATCC CRL1442); human lung cells (W138, ATCC CCL75); human liver cells (HepG2, HB8065); mouse mammary tumor (MMT 060652, ATCC CCL51); rat hepatoma cells (Baumann *et al.*, 1980) and TRI cells (Mather *et al.*, 1982).

It will be appreciated that when expressed in mammalian tissue, the recombinant HIV gene products may have higher molecular weights than expected due to glycosylation. It is therefore intended that partially or completely glycosylated forms of HIV preproteins or peptides having molecular weights somewhat different from 160, 120 or 41 kD are within the scope of this invention.

Other preferred expression vectors are those for use in eukaryotic systems. An exemplary eukaryotic expression system is that employing vaccinia virus, which is well-known in the art. See, for example, Macket *et al.* (1984); Glover, *supra*; and WO 86/07593. Yeast expression vectors are known in the art. See, for example, U.S. Patents 4,446,235; 4,443,539; 4,430,428; and EP 103409; EP 100561; EP 96491.

Another preferred expression system is vector pHSI, which transforms Chinese hamster ovary cells (see WO 87/02062). Mammalian tissue may be cotransformed with DNA encoding a selectable marker such as dihydrofolate reductase (DHFR) or thymidine kinase and DNA encoding the HIV protein or peptide. If wild type DHFR gene is

employed, it is preferable to select a host cell which is deficient in DHFR, thus permitting the use of the DHFR coding sequence as marker for successful transfection in hgt

medium, which lacks hypoxanthine, glycine, and thymidine. An appropriate host cell in this case is the Chinese hamster ovary (CHO) cell line deficient in DHFR activity,

5 prepared and propagated as described by Urlaub & Chasin, (1980).

Depending on the expression system and host selected, HIV proteins or peptides are produced by growing host cells transformed by an exogenous or heterologous DNA construct, such as an expression vector described above under conditions whereby the HIV protein is expressed. The HIV protein or peptide is then isolated from the host cells
10 and purified. If the expression system secretes the protein or peptide into the growth media, the protein can be purified directly from cell-free media. The selection of the appropriate growth conditions and initial crude recovery methods are within the skill of the art.

Once a coding sequence for an HIV protein or peptide of the invention has been
15 prepared or isolated, it can be cloned into any suitable vector and thereby maintained in a composition of cells which is substantially free of cells that do not contain an HIV coding sequence. Numerous cloning vectors are known to those of skill in the art. Examples of recombinant DNA vectors for cloning and host cells which they can transform include the various bacteriophage lambda vectors (*E. coli*), pBR322 (*E. coli*), pACYC177 (*E. coli*),
20 pKT230 (gram-negative bacteria), pGV1106 (gram-negative bacteria), pLAFRI (gram-negative bacteria), pME290 (*non-E. coli* gram-negative bacteria), pHV14 (*E. coli* and *Bacillus subtilis*), pBD9 (*Bacillus*), pIJ61 (*Streptomyces*). pUC6 (*Streptomyces*), actinophage, fC31 (*Streptomyces*). YIpS (*Saccharomyces*), YCp19 (*Saccharomyces*), and bovine papilloma virus (mammalian cells). See generally, Glover, *supra*; T. Maniatis *et*
25 *al.*, *supra*; and Perbal, *supra*.

Fusion Proteins

HIV envelope fusion proteins and methods for making such proteins have been previously described (U.S. Patent 5,885,580). It is now a relatively straight forward
30 technology to prepare cells expressing a foreign gene. Such cells act as hosts and may include, for the fusion proteins of the present invention, yeasts, fungi, insect cells, plants

cells or animals cells. Expression vectors for many of these host cells have been isolated and characterized, and are used as starting materials in the construction, through conventional recombinant DNA techniques, of vectors having a foreign DNA insert of interest. Any DNA is foreign if it does not naturally derive from the host cells used to
5 express the DNA insert. The foreign DNA insert may be expressed on extrachromosomal plasmids or after integration in whole or in part in the host cell chromosome(s), or may actually exist in the host cell as a combination of more than one molecular form. The choice of host cell and expression vector for the expression of a desired foreign DNA largely depends on availability of the host cell and how fastidious it is, whether the host
10 cell will support the replication of the expression vector, and other factors readily appreciated by those of ordinary skill in the art.

The foreign DNA insert of interest comprises any DNA sequence coding for fusion proteins including any synthetic sequence with this coding capacity or any such cloned sequence or combination thereof. For example, fusion proteins coded and
15 expressed by an entirely recombinant DNA sequence is encompassed by this invention but not to the exclusion of fusion proteins peptides obtained by other techniques.

Vectors useful for constructing eukaryotic expression systems for the production of fusion proteins comprise the fusion protein's DNA sequence, operatively linked thereto with appropriate transcriptional activation DNA sequences, such as a promoter and/or
20 operator. Other typical features may include appropriate ribosome binding sites, termination codons, enhancers, terminators, or replicon elements. These additional features can be inserted into the vector at the appropriate site or sites by conventional splicing techniques such as restriction endonuclease digestion and ligation.

Yeast expression systems, which are the preferred variety of recombinant
25 eukaryotic expression system, generally employ *Saccharomyces cerevisiae* as the species of choice for expressing recombinant proteins. Other species of the genus *Saccharomyces* are suitable for recombinant yeast expression system, and include but are not limited to *carlsbergensis*, *uvarum*, *rouxii*, *montanus*, *kluyveri*, *elongisporus*, *norbensis*, *oviformis*, and *diastaticus*. *Saccharomyces cerevisiae* and similar yeasts possess well known
30 promoters useful in the construction of expression systems active in yeast, including but not limited to GAP, GAL10, ADH2, PHO5, and alpha mating factor.

Yeast vectors useful for constructing recombinant yeast expression systems for expressing fusion proteins include, but are not limited to, shuttle vectors, cosmid plasmids, chimeric plasmids, and those having sequences derived from two micron circle plasmids. Insertion of the appropriate DNA sequence coding for fusion proteins into these vectors will, in principle, result in a useful recombinant yeast expression system for fusion proteins where the modified vector is inserted into the appropriate host cell, by transformation or other means. Recombinant mammalian expression system are another means of producing the fusion proteins for the vaccines/immunogens of this invention. In general, a host mammalian cell can be any cell that has been efficiently cloned in cell culture. However, it is apparent to those skilled in the art that mammalian expression options can be extended to include organ culture and transgenic animals. Host mammalian cells useful for the purpose of constructing a recombinant mammalian expression system include, but are not limited to, Vero cells, NIH3T3, GH3, COS, murine C127 or mouse L cells. Mammalian expression vectors can be based on virus vectors, plasmid vectors which may have SV40, BPV or other viral replicons, or vectors without a replicon for animal cells. Detailed discussions on mammalian expression vectors can be found in the treatises of Glover, *DNA Cloning: A Practical Approach*, Vols. 1-4 (Oxford, IRL Press, 1985).

Fusion proteins may possess additional and desirable structural modifications not shared with the same organically synthesized peptide, such as adenylation, carboxylation, glycosylation, hydroxylation, methylation, phosphorylation or myristylation. These added features may be chosen or preferred as the case may be, by the appropriate choice of recombinant expression system. On the other hand, fusion proteins may have its sequence extended by the principles and practice of organic synthesis.

25

Vaccines and Immunogenic Compositions

When used in vaccine or immunogenic compositions, the proteins or peptides of the present invention may be used as "subunit" vaccines or immunogens. Such vaccines or immunogens offer significant advantages over traditional vaccines in terms of safety and cost of production; however, subunit vaccines are often less immunogenic than

30

whole-virus vaccines, and it is possible that adjuvants with significant immunostimulatory capabilities may be required in order to reach their full potential.

Currently, adjuvants approved for human use in the United States include aluminum salts (alum). These adjuvants have been useful for some vaccines including hepatitis B, diphtheria, polio, rabies, and influenza. Other useful adjuvants include Complete Freund's Adjuvant (CFA), Incomplete Freund's Adjuvant (IFA), Muramyl dipeptide (MDP) (see Ellouz *et al.*, 1974), synthetic analogues of MDP (reviewed in Chedid *et al.*, 1978), N-acetylmuramyl-L-alanyl-D-isoglutamyl-L-alanine-2-[1,2-dipalmitoyl-s-glycero-3-(hydroxyphosphoryloxy)]ethylamide (MTP-PE) and compositions containing a metabolizable oil and an emulsifying agent, wherein the oil and emulsifying agent are present in the form of an oil-in-water emulsion having oil droplets substantially all of which are less than one micron in diameter (see EP 0399843).

The formulation of a vaccine or immunogenic compositions of the invention will employ an effective amount of the protein or peptide antigen. That is, there will be included an amount of antigen which, in combination with the adjuvant, will cause the subject to produce a specific and sufficient immunological response so as to impart protection to the subject from subsequent exposure to an HIV virus. When used as an immunogenic composition, the formulation will contain an amount of antigen which, in combination with the adjuvant, will cause the subject to produce specific antibodies which may be used for diagnostic or therapeutic purposes.

The vaccine compositions of the invention may be useful for the prevention or therapy of HIV-1 infection. While all animals that can be afflicted with HIV-1 can be treated in this manner, the invention, of course, is particularly directed to the preventive and therapeutic use of the vaccines of the invention in man. Often, more than one administration may be required to bring about the desired prophylactic or therapeutic effect; the exact protocol (dosage and frequency) can be established by standard clinical procedures.

The vaccine compositions are administered in any conventional manner which will introduce the vaccine into the animal, usually by injection. For oral administration the vaccine composition can be administered in a form similar to those used for the oral administration of other proteinaceous materials. As discussed above, the precise amounts

and formulations for use in either prevention or therapy can vary depending on the circumstances of the inherent purity and activity of the antigen, any additional ingredients or carriers, the method of administration and the like.

By way of non-limiting illustration, the vaccine dosages administered will typically be, with respect to the gp120 antigen, a minimum of about 0.1 mg/dose, more typically a minimum of about 1 mg/dose, and often a minimum of about 10 mg/dose. The maximum dosages are typically not as critical. Usually, however, the dosage will be no more than 500 mg/dose, often no more than 250 mg/dose. These dosages can be suspended in any appropriate pharmaceutical vehicle or carrier in sufficient volume to carry the dosage. Generally, the final volume, including carriers, adjuvants, and the like, typically will be at least 0.1 ml, more typically at least about 0.2 ml. The upper limit is governed by the practicality of the amount to be administered, generally no more than about 0.5 ml to about 1.0 ml.

Peptides of the invention corresponding to domains of the envelope protein such as V3 may be constructed or formulated into compounds or compositions comprising multimers of the same domain or multimers of different domains. For instance, peptides corresponding to the V3 domain may be circularized by oxidation of the cysteine residues to form multimers containing 1, 2, 3, 4 or more individual peptide epitopes. The circularized form may be obtained by oxidizing the cysteine residues to form disulfide bonds by standard oxidation procedures such as air oxidation.

Synthesized peptides of the invention may also be circularized in order to mimic the geometry of those portions as they occur in the envelope protein. Circularization may be facilitated by disulfide bridges between existing cysteine residues. Cysteine residues may also be included in positions on the peptide which flank the portions of the peptide which are derived from the envelope protein. Alternatively, cysteine residues within the portion of a peptide derived from the envelope protein may be deleted and/or conservatively substituted to eliminate the formation of disulfide bridges involving such residues. Other means of circularizing peptides are also well known. The peptides may be circularized by means of covalent bonds, such as amide bonds, between amino acid residues of the peptide such as those at or near the amino and carboxy termini (see U.S. Patent 4,683,136).

In an alternative format, vaccine or immunogenic compositions may be prepared as vaccine vectors which express the HIV protein or peptide of the invention in the host animal. Any available vaccine vector may be used, including live Venezuelan Equine Encephalitis virus (see U.S. Patent 5,643,576), poliovirus (see U.S. Patent 5,639,649), pox virus (see U.S. Patent 5,770,211) and vaccina virus (see U.S. Patents 4,603,112 and 5,762,938). Alternatively, naked nucleic acid encoding a protein or peptide of the invention may be administered directly to effect expression of the antigen (see U.S. Patent 5,739,118).

10 *Diagnostic Reagents*

The HIV protein or peptide compositions of the present invention may be used as diagnostic reagents in immunoassays to detect anti-HIV antibodies, particularly anti-gp120 antibodies. Many HIV immunoassay formats are available. Thus, the following discussion is only illustrative, not inclusive. See generally, however, U.S. Patents 4,743,678; 4,661,445; and 4,753,873 and EP 0161150 and EP 0216191.

Immunoassay protocols may be based, for example, upon composition, direct reaction, or sandwich-type assays. Protocols may also, for example, be heterogeneous and use solid supports, or may be homogeneous and involve immune reactions in solution. Most assays involved the use of labeled antibody or polypeptide. The labels may be, for example, fluorescent, chemiluminescent, radioactive, or dye molecules. Assays which amplify the signals from the probe are also known, examples of such assays are those which utilize biotin and avidin, and enzyme-labeled and mediated immunoassays, such as ELISA assays.

Typically, an immunoassay for anti-HIV antibody will involve selecting and preparing the test sample, such as a biological sample, and then incubating it with an HIV protein or peptide composition of the present invention under conditions that allow antigen-antibody complexes to form. Such conditions are well known in the art. In a heterogeneous format, the protein or peptide is bound to a solid support to facilitate separation of the sample from the polypeptide after incubation. Examples of solid supports that can be used are nitrocellulose, in membrane or microtiter well form, polyvinylchloride, in sheets or microtiter wells, polystyrene latex, in beads or microtiter

plates, polyvinylidene fluoride, diazotized paper, nylon membranes, activated beads, and Protein A beads. Most preferably, Dynatech, Immulon® microtiter plates or 0.25 inch polystyrene beads are used in the heterogeneous format. The solid support is typically washed after separating it from the test sample.

5 In homogeneous format, on the other hand, the test sample is incubated with the HIV protein or peptide in solution, under conditions that will precipitate any antigen-antibody complexes that are formed, as is known in the art. The precipitated complexes are then separated from the test sample, for example, by centrifugation. The complexes formed comprising anti-HIV antibody are then detected by any number of
10 techniques. Depending on the format, the complexes can be detected with labeled anti-xenogenic Ig or, if a competitive format is used, by measuring the amount of bound, labeled competing antibody. These and other formats are well known in the art.

Diagnostic probes useful in such assays of the invention include antibodies to the HIV-1 envelope protein. The antibodies may be either monoclonal or polyclonal,
15 produced using standard techniques well known in the art (See Harlow & Lane, *Antibodies: A Laboratory Manual*, (Cold Spring Harbor, Cold Spring Harbor Laboratory Press, 1988). They can be used to detect HIV-1 envelope protein by specifically binding to the protein and subsequent detection of the antibody-protein complex by ELISA, Western blot or the like. The HIV-1 envelope protein used to elicit these antibodies can
20 be any of the variants discussed above. Antibodies are also produced from peptide sequences of HIV-1 envelope proteins using standard techniques in the art (Harlow & Lane, *supra*). Fragments of the monoclonals or the polyclonal antisera which contain the immunologically significant portion can also be prepared.

The following working examples specifically point out preferred embodiments of
25 the present invention, and are not to be construed as limiting in any way the remainder of the disclosure. Other generic configurations will be apparent to one skilled in the art. All references, including U.S. or foreign patents, referred to in this application are herein incorporated by reference in their entirety.

30 Examples

The following methods were used in the Examples:

Reference Serum Donor Envelope Gene Cloning

The donor of the HIV-1 Neutralizing Serum (2) (Reference 2), available in the NIH-AIDS-Research-and-Reference-Reagent-Program-(Catalog-Number: 1983) is a participant in a long term cohort study at the Clinical Center, NIH (Vujcic *et al.*, 1995).

- 5 The blood used to prepare Reference 2 had been collected in the Spring of 1989. Peripheral blood mononuclear cells that had been cryopreserved from donations obtained approximately six months and one year prior to the time of Reference 2 collections were used as sources of DNA for *env* gene cloning. The cells had not been stored to maintain viability. DNA was extracted using phenol/chloroform from approximately $1-3 \times 10^6$
- 10 cells from each donation (Quinnan *et al.*, 1998). The DNA was used as template in a nested polymerase chain reaction, similar to that described previously, except rTth was used as the DNA polymerase, following the manufacturer's instructions (Barnes, 1992; Cariello *et al.*, 1991). The DNA was cloned into the expression vector pSV7d, as previously described (Quinnan *et al.*, 1998; Stuve *et al.*, 1987).

15

Other env Gene Clones and Virus Pools

The following HIV-1 *env* clones in the expression vector pSV3 were obtained from the AIDS Research and Reference Reagent Program, 93MW965.26 (clade C), 92RWO20.5 (clade A), 93TH966.8 (clade E), 92UG975.10 (clade G) (Gao *et al.*, 1994).

- 20 The production of *env* clones from the molecular virus clones NL43, AD8, and SF162 has been previously described (Quinnan *et al.*, 1998; Adachi *et al.*, 1986; Theodore *et al.*, 1996; Englund *et al.*, 1995). *env* gene of the Z2Z6 strain was cloned similarly, using molecular virus clone plasmid as template in polymerase chain reaction, and cloning the genes into the plasmid pSV7d (Seth *et al.*, 1993). The production of primary isolate *env*
- 25 clones from participants in the Multicenter AIDS Cohort Study, designated here P9 and P10, has also been previously described (Quinnan *et al.*, 1998). P9 and P10- virus pools were prepared by single subpassages of the cell culture media from primary cultures in PHA blasts (Quinnan *et al.*, 1998). The use of molecular virus clones for preparation of virus pools of NL43 in H9 cells, and NL(SF162) and AD8, in PHA blasts, has also been
- 30 previously described (Quinnan *et al.*, 1998).

Cell Cultures

The H9 cell line was obtained from Robert Gallo (Mann *et al.*, 1989). The Molt 3 cell-line was obtained from the American-Type-Culture Collection, Rockville, MD (ATCC). (Daniel *et al.*, 1988) The HOS cell lines expressing CD4 and various coreceptors for HIV-1 were obtained from the NIH AIDS Research and Reference Reagent Program, as was the PM1 cell line (Deng *et al.*, 1996; Landau *et al.*, 1992; Lusso *et al.*, 1995). The 293T cell line was obtained from the ATCC, with permission from the Rockefeller Institute (Liou *et al.*, 1994). The H9, Molt3 and PM1 cell cultures were maintained in RPMI-1640 medium supplemented with 10% fetal bovine serum and antibiotics (Gibco). The HOS and 293T cells were maintained in Dulbecco's Minimal Essential Medium (Gibco), with similar supplements, except that the HOS cell medium was supplemented with puromycin for maintenance of plasmid stability. Cryopreserved human peripheral blood lymphocytes were stimulated with PHA and used for virus infections (Quinnan *et al.*, 1998; Mascola *et al.*, 1994).

Reverse Transcriptase Assay

Reverse transcriptase activity was assayed as previously described (Park *et al.*, 1998).

Virus Neutralization Assays

The virus NL43 was used in neutralization assays which employed Molt3 cells as target cells and used giant cell formation for endpoint determination, as previously described (Vujcic *et al.*, 1995). The amounts of virus used were sufficient to result in the formation of 30-50 giant cells per well (Vujcic *et al.*, 1995; Lennette, 1964). The viruses, NL(SF162) and AD8, P9 and P10 were tested for neutralization in PHA stimulated human lymphoblasts in the presence of IL-2 (Quinnan *et al.*, 1998; Mascola *et al.*, 1994). In the latter assays ten percent of the cell suspension was removed each week, fifty percent of the medium was changed each week, and medium was sampled twice weekly from each well for reverse transcriptase assay. The reverse transcriptase assays were performed on the test samples from the first sampling date at which the non-neutralized control wells had reverse transcriptase activity about 10-20 x background, generally on

day 14 or 17 of the assay. The neutralization endpoint was considered to be the highest dilution of serum at which reverse transcriptase activity was reduced at least fifty percent.

The Reference Neutralizing Sera 1 and 2 and the Negative Reference Serum were used as positive and negative controls (NIH AIDS Research and Reference Reagent Program)

5

Pseudovirus Construction and Assays of Pseudoviruses for Infectivity and Neutralization

Pseudoviruses were constructed and assayed using methods similar to those described previously (Quinnan *et al.*, 1998; Deng *et al.*, 1996; Park *et al.*, 1998). pSV7d-*env* plasmid DNA and pNL43.luc+.E-R- were cotransfected into 70 to 80% confluent 293T cell cultures using the calcium phosphate/Hepes buffer technique, following manufacturer's instructions (Promega, Madison, WI), in 24 well plastic tissue culture trays or 25 cm² flasks (Quinnan *et al.*, 1998; Deng *et al.*, 1996; Park *et al.*, 1998). After 24 hours the medium was replaced with medium containing one mM sodium butyrate (Quinnan *et al.*, 1998; Park *et al.*, 1998). Two days after transfection medium was harvested, passed through a 45µm sterile filter (Millipore Corp, Bedford, MA), supplemented with an additional 20% fetal bovine serum and stored at -80°C.

Pseudovirus infectivity was assayed in PM1 or HOS-CD4 cells expressing various co-receptors. Transfection supernatants were serially diluted and inoculated into cells in 96 well plates, 50 µl per well. Assays were routinely performed in triplicate. The cultures were incubated for four days, centrifuged at 400 x g for ten minutes if PM1 cells were used, and medium removed by aspiration. The cells were washed twice with phosphate buffered saline, lysed with 25 µl cell culture lysing reagent according to the manufacturer's instructions (Promega, Madison, WI); the cells were then tritated into the medium, and 10 µl of the suspensions were transferred to wells of 96 well luminometer plates. Substrate was added in 100 µl volumes automatically, and the luminescence read using a MicroLumatPlus luminometer (EG&G Berthold, Hercules, CA). Mock PV controls were used in each assay consisting of media harvested from 293T cell cultures transfected with pSV7d (without an *env* insert) and pNL43.Luc.E-R-, and processed in the same way as cultures for PV preparation. Infectivity endpoints were determined by a modified Reed Munch method; an individual well was considered positive if the luminescence was at least 10-fold greater than the mock control, and the endpoint was

considered to be the highest dilution at which the calculated frequency of positivity was \geq 50% (Quinnan *et al.*, 1998; Park *et al.*, 1998; Lennette, 1964). Luminescence resulting from infection with minimally diluted samples was generally about 10,000-fold greater than background.

5 Neutralization tests were performed using PM1 or HOS-CD4 cells. Aliquots of 25 μ l of two-fold serial serum dilutions were mixed with equal volumes of diluted PV in wells of 96 well plates. The PV dilutions were selected so as to expect luminescence in the presence of non-neutralizing serum of about 100-fold of background. Assays were performed in triplicate. The virus serum mixtures were incubated for sixty minutes at 10 40°C, after which 150 μ l aliquots of PM1 cell suspensions were added, which each contained 1.5×10^4 cells, or the suspensions were transferred to wells containing HOS-CD4 cells. The assays were then processed similarly to the infectivity assays. The neutralization endpoints were calculated by a modified Reed-Munch method in which the endpoint was considered to be the highest serum dilution calculated to have a frequency 15 of \geq 50% for reducing luminescence by \geq 90% compared to the non-neutralized control. PV titrations were conducted in duplicate in parallel with each neutralization assay.

Nucleic Acid Sequencing

Nucleotide sequence analysis was performed using the di-deoxy cycle sequencing 20 technique and AmpliTaq FS DNA polymerase, according to manufacturer's directions (Perkin Elmer Applied Biosystems, Foster City, CA). After the sequencing reaction the DNA was purified using Centriflex Gel Filtration Cartridges (Advanced Genetic Technologies, Gaithersburg, MD). Sequencing gels were run and analyzed using an Applied Biosystems Prism, Model 377 DNA Sequencer. Sequencing was performed on 25 both strands. Sequence alignment was performed using the Editseq SEQMAN, and Megalign programs in DNA Star according to the method of Higgins and Sharp (1989).

Example 1: Comparability of Clones Isolated from Different Time Points

From the samples of patient cells from each of the two time points, *env* clones 30 were recovered which encoded proteins which were capable of mediating pseudovirus entry into target cells. Two such clones from each time point were further characterized.

As shown in Table 1, the envelopes of all four clones mediated infection for PM1 cells and were neutralized comparably by References 1 and 2. Pseudoviruses carrying envelopes corresponding to each clone were also tested for infectivity for HOS-CD4 cells expressing either CXCR4 or CCR5, and all four were infectious only for the cells expressing CCR5, as shown in Table 2. Nucleotide sequences including the V3 regions were analyzed for each clone, with more than 300 bases assigned for each, and no differences between the clones were found (results not shown). Based on the absence of demonstration of differences in these assays, a single clone from the March sample was selected for use in subsequent assays, and is designated R2, hereafter.

Example 2: Clone R2 Genotype and Host Range Phenotype

The complete nucleotide sequence of the *env* gene clone R2 was determined, and found to have an open reading frame of 2598 bases (Genbank Accession Number: AF128126). The amino acid sequence deduced from this sequence is shown in Table 3 (SEQ ID NO:1). There are thirty predicted glycosylation sites, compared to twenty-nine in the consensus clade B sequence; four consensus glycosylation sites are lacking in R2, including those at residues 146, 215, 270, and 368 (numbering according to the Human Retroviruses and AIDS Database clade B consensus sequence), in the V2, C2, C2 and V4 regions of gp120, respectively (Myers *et al.*, 1993). The consensus glycosylation sequences at residues 215 and 270 are highly and moderately variable, respectively.

Genotypic analyses conducted included evaluation of the gp120 and gp41 nucleotide coding sequences in comparison to those of a number of strains of clades A through G, as shown in Figure 1 (Saitou *et al.*, 1987; Myers & Miller, 1988). Both coding regions were more closely related to clade B than non-clade B sequences. Comparative analyses of regions of the predicted gp120 and gp41 amino acid sequences were also performed (results not shown). The regions analyzed included: each constant and variable region of gp120; the proximal gp41 ectodomain including the leucine zipper region; the part of gp41 extending from the end of the leucine zipper to the second cysteine; the remaining gp41 ectodomain, and the transmembrane region; and the cytoplasmic region. R2 consistently related more closely with the clade B sequences than the others.

Example 3: Comparative Sensitivity of R2 and Other Clade B Viruses and Pseudoviruses to Neutralization by Sera from Individuals with Clade B Infections

The neutralization of R2 pseudovirus was compared to other clade B viruses and
 5 pseudoviruses as shown in Figure 2. Of the five virus-pseudovirus comparisons made
 (P9, P10, NL43, AD8 and SF162 V and PV), there were no significant differences in the
 neutralization of matched viruses and pseudoviruses by paired t test (statistical results not
 shown). Each of the pseudovirus preparations was neutralized by seven, eight, or nine of
 the sera tested, and the geometric mean titers ranged from 1:13.9 to 1:56, while the R2-
 10 PV was neutralized by all ten of the sera tested, with a geometric mean titer of 1:73.5.
 Although the neutralization titers of each of the different sera against R2 and the other
 pseudoviruses were frequently within four-fold, the neutralization of R2-PV was
 significantly greater by paired t test than four of the other PV preparations.

15 *Example 4: Comparative neutralization of pseudoviruses expressing R2 and other envelopes of diverse subtypes by sera from diverse subtype infections.*

The results of comparative neutralization testing using sera from individuals
 infected with HIV- I strains of subtypes A, C and E, and the Reference 1 and 2, and one
 Thai clade B serum are shown in Table 4. Reference 2 neutralized the pseudovirus
 20 expressing the homologous R2 envelope at the modest titer of 1:64 in the experiment
 shown and within two-fold of this titer in many other experiments. It neutralized the
 other seven pseudoviruses tested at low to moderate titers, as well. The R2 pseudovirus
 was neutralized by seventeen of twenty-four sera, including sera from people infected
 with each of the clades A-F. The other two clade B pseudoviruses were neutralized less
 25 frequently and were also neutralized infrequently by the clade E sera. The frequency of
 neutralization by sera from individuals infected with different clades was not significantly
 skewed for any of the other four pseudoviruses. Clade A, C, D and G pseudoviruses were
 neutralized by eight, seventeen, six and three of the seventeen sera tested, respectively.
 The clade C pseudovirus was substantially more sensitive to neutralization, in general
 30 than the others tested. The clade E pseudovirus was neutralized by five of five clade D

sera and seven of eight clade E sera but only one of the sera from people infected by other clades.

Example 5: Synthetic peptides generated from V3 amino acid sequences from R2 strain.

5 R2 strain V3 peptides were synthesized using an automated ABI synthesizer and Fmoc chemistry (Zeng *et al.*, 1997). The sequences of these peptides were KSIPMGPGRAFYTGTGQI (SEQ ID NO:2) and CSRPNNTRKSIPMGPGRAFYTGTGQIIGDIRQAH (SEQ ID NO:3). The mutant R2(313-4PM/HL, 325Q/D) V3 peptide was prepared similarly. Strain 93TH966.8 V3
10 peptide, sequence: CTRPSNNTRTSTTIGPGQVFYRTGDITGNIRKAYC (SEQ ID NO:4) was synthesized using the same methods. The peptides were purified using C18, acetonitrile-in-water gradient chromatography with a Waters High Performance Liquid Chromatograph. Sequences of the purified peptides were verified using an ABI automated sequencer. Peptides were lyophilized and stored at 4-8°C. Preparation of a
15 linear MN strain V3 peptide has been described previously (Carrow *et al.*, 1991). Cyclic MN strain 35-mer peptide was obtained from the AIDS Research and Reference Reagent Program (Catalog #1841) provided by Catasti *et al.*, (1996).

The R2 V3 35-mer was insoluble in water, while all other peptides tested were soluble in water to at least 10 mg/ml. To obtain cyclic peptides, solutions of the R2 and
20 R2(313-4PM/HL, 325Q/D) V3 35-mers in dimethylsulfoxide (DMSO), 10 mg/ml, were diluted 1:10 in water at room temperature or 37°C and the pH was adjusted to 8.5 with ammonium hydroxide. These solutions were aerated by bubbling air through the solutions for periods ≥ 1 hour. Following aeration, the pH was adjusted to 7.4 using hydrochloric acid. A portion of the R2 35-mer peptide precipitated during these
25 procedures. To obtain an approximate quantitation of the amount of R2 V3 35-mer that remained in solution, the turbidity of the suspension was determined at 480 nm wavelength visible light using a spectrophotometer. The spectrophotometer was blanked with a solution of 10 percent DMSO in water, and a standard curve was produced using slurries of known amounts of the 35-mer peptide suspended in water. The amount of
30 precipitate estimated by turbidity was subtracted from the amount of peptide added at the beginning of the preparation procedure to estimate the amount remaining in solution. The

solubility of the oxidized R2 35-mer peptide in 10 percent DMSO solution at pH=7.4 was estimated to be 300-350 $\mu\text{g/ml}$ when processed at room temperature, or 850-900 $\mu\text{g/ml}$ when processed at 37°C. Peptides were sterilized by passage through 0.22 μ pore size filters prior to use.

5

Example 6: Peptide blocking of neutralizing antibody activity against clone R2 pseudovirus.

The neutralization blocking effects of synthetic V3 peptides were examined to test the contribution of V3-anti-V3 interactions in the neutralizing cross reactivities of Reference 2 and clone R2. The blocking effects of peptides on neutralizing activity of Reference 2 against clone R2 pseudovirus are shown in Figure 3A. Usually, the linear 17-mer peptide had no inhibitory effect on neutralization, as shown. In only one of several experiments two-fold reduction of neutralization was observed in the presence of 17-mer peptide. Concentration-dependent inhibitory effects of the cyclic 35-mer R2 V3 peptide on neutralization of clone R2 pseudovirus by Reference 2 was observed in the experiment shown and in numerous other similar experiments. Maximum effect was observed at approximately 15 $\mu\text{g/ml}$. No inhibitory effect was observed using a cyclic peptide homologous to the V3 region of the HIV-1 93TH966.8 strain.

The comparative effects of the R2 and MN strain V3 peptides on neutralization of the R2 and MN strain pseudoviruses are shown in Figure 3B. The results shown are representative of two additional experiments. Only the cyclic R2 V3 peptide produced consistent blocking of R2 pseudovirus neutralization. The linear R2 and MN, and the cyclic MN peptides did not block R2 neutralization in two experiments and blocked only two-fold in a third experiment. In contrast, the MN cyclic and linear peptides consistently inhibited MN strain neutralization eight- to sixteen-fold in these experiments, and the R2 peptides had consistent two-fold inhibitory effects on neutralization of the MN strain. These effects of MN peptides on MN strain neutralization are consistent with previous reports (Carrow *et al.*, 1991; Park *et al.*, 1999).

Example 7: Cyclic R2 V3 peptide inhibition of neutralization of R2 pseudoviruses by sera from MACS patients.

Inhibition of heterologous serum neutralization of R2 pseudovirus by cyclic R2 V3 peptide was evaluated to determine if cross reactivity of these sera with R2 included effects of anti-V3 antibodies. The comparative neutralization titers of sera from ten patients from the MACS against clone R2 pseudovirus in the presence and absence of cyclic R2 V3 peptide are shown in Figure 4A (Quinnan *et al.*, 1998). These sera have been described previously, and have been shown to neutralize primary HIV-1 enveloped pseudoviruses cross reactively, but to a lesser extent than Reference 2 (Zhang *et al.*, 1999). Each serum was tested twice. Seven of the sera appeared to be inhibited at least two-fold in one or both experiments. The geometric mean inhibitory effect of all the tests was 1.9-fold. The results of twelve tests conducted at the same times as those tests shown in Figures 4A and 4B are shown for Reference 2; the geometric mean inhibitory effect was 3.56.

Example 8: Cyclic R2 V3 peptide inhibition of Reference 2 neutralization of pseudoviruses expressing envelopes from the MACS patients.

Inhibition of Reference 2 neutralization of pseudoviruses expressing heterologous envelopes by cyclic R2 V3 peptide was evaluated to determine whether anti-V3 antibody contributed to the neutralizing cross reactivity of Reference 2. The results of these experiments are shown in Figure 4B. Each pseudovirus was tested two or three times. The peptide appeared to exert a two-fold inhibitory effect in one, two, or three of the experiments using each of the six pseudoviruses. The geometric mean inhibitory effect was 1.6-fold.

Example 9: Induction of cross-reactive neutralizing antibodies in mice following immunization with recombinant delivery vectors encoding HIV-1 envelope proteins.

The DNA clone encoding the R2 envelope was introduced into an expression vector which can be used to express the envelope protein complex *in vivo* for immunization. The recombinant delivery vector expressing the R2 envelope clone was been administered to mice, both in its full length, encoding both gp120 and gp41, or in a truncated form. The truncated form is secreted by cells which express gp140. Both the full-length and truncated form of these constructs induced neutralizing antibodies in mice.

The mice which received the gp140 construct, which includes the V3 region, have developed neutralizing antibodies which neutralize at least three different strains of HIV-1, including the R2-strain, a macrophage-tropic laboratory strain known as SF162,

- and a primary strain which is not laboratory adapted. The amount of cross-reactivity
5 observed exceeds that induced by most or all other HIV immunogens that have been tested as single agents.

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2

From Donor 2

Serum	Neutralization Titer Against Clone			
	10.1	10.2	3.1	3.2
Reference 1	1:32	1:64	1:32	1:64
Reference 2	1:128	1:128	1:128	1:128

Table 2. Coreceptor Dependency of R2 Pseudovirus Entry Into HOS-CD4 Cells

Pseudo-virus	Infectivity-Titer						In PM1 Cells
	In HOS-CD4 Cells Expressing						
	CCR1	CCR2b	CCR3	CCR4	CCR5	CXCR4	
R2	<1:4	<1:4	<1:4	<1:4	1:64	<1:4	1:32
P9	<1:4	<1:4	<1:4	<1:4	1:256	<1:4	1:8
NL4-3	<1:4	<1:4	<1:4	<1:4	1:32	>1:256	1:8
AD8	<1:4	<1:4	<1:4	<1:4	1:256	<1:4	1:32

Table 3. Inferred Amino Acid Sequence of the R2 Envelope Clone from Donor 2.

Amino Acid Residue ^a	Residue Number
MRVKGIRRNY QHWWGWGTML LGLLMICSAT EKLWVTVYYG VPVWKEATTT	50
LFCASDAKAY DTEAHNVWAT HACVPTDPNP QEVELVNVTE NFNMWKNNMV	100
EQMHEDIISL WDQSLKPCVK LTPLCVTLNC TDLRNTTNTN N STDNNNSNS	150
EGTIKGGEMK NCSFNIAISI GDKMQKEYAL LYKLDIEPID NDNTSYRLIS	200
CNTSVITQAC PKISFEPIPI HYCAPAGFAI LKCNDKKFSG KGSCKNVSTV	250
QCTHGIRPVV STQLLNGSL AEEEVVIRSE NFTNNAKTII VQLREPVKIN	300
CSRPNNNTRK SIPMGPGRAF YTTGQIIGDI RQAHCNISKI N WTNALKQVV	350
EKLGEQFNKT KIVFTNSSGG DPEIVTHSFN CAGEFFYCNT TQLFDSIWNS	400
ENGTWNITRG LNNTGRNDTI TLPCRKIQII NRWQEVGKAM YAPPIKGNIS	450
CSSNITGLLL TRDGGKDDNS RDGNETFRPG GGDMRDNWRN ELYKYKVVKI	500
EPLGVAPTKA KRRVVQREER AVGLGAMFIG FLGAAGSTMG AASVTLTVQA	550
RQLLSGIVQQ QSNLLRAIEA QHLLQLTVW GIKQLQARIL AVERYLKDQQ	600
LLGIWGCSSK LICTTTVPWN ASWSKNKTLE AIWNNMTWMQ WDKEIDNYTK	650
LIYSLIEESQ IQQEKNEQEL LELDKWANLW NWFDISNWLW YIKIFIMIVG	700
GLVGLRIVFV VLSIVNRVRQ GYSPLSFQTR LPAPRGPDRP EEIEEEGGDR	750
DRDRSGLLVD GFLTIIWVDL RSLCLFSYHR LRDLLLVTR IVELLGRRGW	800
EILKYWWNLL QYWSQELKNS AVSLFNATAI AVAEGTDRVI EVLQRVGRAL	850
LHIPTRIRQG LERALL	866

^aAmino acid residues are identified by standard single letter designations. Predicted N-linked glycosylation sites are indicated by shading and bolding.

Table 4. Neutralization of Pseudoviruses Expressing Envelopes of Various Clades by Sera from People Infected with Various Clades of HIV-1

Clade	Serum ^b	NA Titer Against Pseudovirus (Clade) ^a							
		R2 (B)	P9 (B)	P10 (B)	BR020 (A)	MW965 (C)	Z2Z6 (D)	TH966 (E)	UG975 (G)
B	Ref 1	32	16	32	<10	256	10	<8	<10
	Ref 2	64	32	64	10	128	40	8	10
	WR8465	20	NT ^c	80	<10	640	10	<10	10
A	37570	320	160	20	80	2560	<10	<10	<10
	35374	40	<10	<10	<10	640	<10	<10	<10
	35837	40	20	<10	80	2560	<10	<10	<10
C	5107	40	10	<10	10	1280	<10	<10	<10
	5708	10	<10	<10	<10	320	<10	<10	<10
	5218	80	<10	<10	<10	1280	<10	<10	<10
D	UG9240	<10	NT	NT	NT	NT	NT	20	NT
	UG9370	<10	NT	NT	NT	NT	NT	10	NT
	UG9386	<10	NT	NT	NT	NT	NT	10	NT
	UG93097	10	NT	NT	NT	NT	NT	10	NT
	UG94118	10	NT	NT	NT	NT	NT	20	NT
E	WR5659	10	<10	<10	<10	20	<10	40	<10
	WR5901	<10	<10	<10	40	320	10	40	10
	WR8177	<10	<10	<10	40	640	10	80	<10
	WR8657	<10	<10	10	10	640	<10	80	<10
	WR8593	<10	<10	<10	<10	160	10	40	<10
	1008	<10	<10	<10	<10	10	<10	<10	<10
	1053	20	<10	<10	<10	40	<10	20	<10
	1062	20	10	<10	10	320	<10	20	<10

F	BR9318	<10	NT	NT	NT	NT	NT	<10	NT
	BR93019	10	NT	NT	NT	NT	NT	<10	NT
	BR93020	20	NT	NT	NT	NT	NT	<10	NT
	BR93029	10	NT	NT	NT	NT	NT	<10	NT

^bSera were the Reference Neutralizing Human Serum 1 and 2, or were provided by Dr. J. Mascola, HIVNET, or the UNAIDS Program, as described in the text.

NT=not tested.

Sub. 23
 Table 5. Comparison of V3 Region Amino Acid Sequences of Clone R2 with Phenetic Subgroup Consensus Sequences 1 Through 13 and Clade A Through E Consensus Sequences.*

Clone, Subgroup or Clade	V3 Region Amino Acid Sequence
R2	NNTR.KSIPMGPGRAFYTTGQIIGDIRQAHC
PHENETIC 1	----.---HI-----D-----
PHENETIC 2	----.---SI-----A-E-----
PHENETIC 3	----.---SI-----A-K-----
PHENETIC 4	----.---RI---Q---A-D-----
PHENETIC 5	----.---HI-----A-K-----
PHENETIC 6	K--RRR-H.I-----K-----
PHENETIC 7	----.T--TI---QV--R--K-----
PHENETIC 8	KKM-.T-ARI---V-HK--D---S-TK-Y-
PHENETIC 9	----.Q-THI---Q-L---.D---K-----
PHENETIC 10	----.QGTHI-----Y---.N-----
PHENETIC 11	----.QRTSI-Q-QAL---.E-R-----A-
PHENETIC 12	D-IKIQRT-I-Q-Q-L---RITGYI.G----
PHENETIC 13	Q-K-.QGT-I-L-Q-L---R.-K---K---
CLADE A	----.---VHI---Q---A-D-----
CLADE B	----.---HI-----E-----
CLADE C	----.---RI---QT-YA--D-----
CLADE D	----.QRTHI---Q-L---.R-----
CLADE E	----.T--TI---QV--R--D-----K-Y-

*Dashes indicate residues at which the individual sequences are identical to R2. The periods indicate sites of insertions or deletions.

References

- Adachi A, Gendelman HE, Keonig S, Folks T, Wiley R, Rabson A and Martin MA, *Production of acquired immunodeficiency syndrome-associated retrovirus in human and nonhuman cells transfected with an infectious molecular clone*. J Virol. (1986) 59: 284-291.
- Back NK, Smit L, Schutten M, Nara PL, Tersmette M and Goudsmit J, *Mutations in human immunodeficiency virus type 1 gp41 affect sensitivity to neutralization by gp120 antibodies*. J Virol. (1993) 67: 6897-6902.
- Barnes WM, *The fidelity of Taq polymerase catalyzing PCR is improved by an N-terminal deletion*. Gene (1992) 112: 29-35.
- Baumann H, Gelehrter TD and Doyle D, *Dexamethasone regulates the program of secretory glycoprotein synthesis in hepatoma tissue culture cells*. J Cell Biol (1980) 85: 1-8.
- Cariello NF, Swenberg JA and Skopek TR, *Fidelity of Thermococcus litoralis DNA polymerase (Vent) in PCR determined by denaturing gradient gel electrophoresis*. Nucleic Acids Res. (1991) 19: 4193-4198.
- Catasti P, Bradbury EM and Gupta G. *Structure and polymorphism of HIV-1 third variable loops*. J Biol Chem. (1996) 271: 8236-8242.
- Carrow EW, Vujcic LK, Glass WL, Seamon KB, Rastogi SC, Hendry RM, Boulos R, Nzila N and Quinnan GV. *High prevalence of antibodies to the gp120 V3 region principal neutralizing determinant of HIV-1 MN in sera from Africa and the Americas*. AIDS Res Hum Retroviruses (1991) 7: 831-838.
- Chackerian B, Rudensey LM and Overbaugh J. *Specific N-linked and O-linked glycosylation modifications in the envelope V1 domain of simian immunodeficiency virus variants that evolve in the host alter recognition by neutralizing antibodies*, J Virol (1997) 71: 7719-7727.
- Chedid L, Audibert F and Johnson AG, *Biological activities of muramyl dipeptide, a synthetic glycopeptide analogous to bacterial immunoregulating agents*. Prog Allergy (1978) 25: 63-105.

Daniel MD, Li Y, Naidu YM, Durda PJ, Schmidt DK, Troup CD, Silva DP, MacKey JJ, Kestler HW, Sehgal PK *et al.*, *Simian immunodeficiency virus from African green monkeys*. J Virol. (1988) 62: 4123-4128.

- Deng H, Liu R, Ellmeier W, Choe S, Unutmaz D, Burkhart M, Di Marzio P,
 5 Marmon S, Suttom RE, Hill CM, Davis CB, Peiper SC, Schall TJ, Littman DR and Landau NL, *Identification of a major co-receptor for primary isolates of HIV*. Nature (1996) 381: 661-666.

- D'Souza MP, Durda P, Hanson CV, Milman G and collaborating investigators, *Evaluation of monoclonal antibodies to HIV-1 by neutralization and serological assays: an international collaboration*. AIDS (1991) 5: 1061-1070.

Ellouz F, Adam A, Ciorbaru R and Lederer E, *Minimal structural requirements for adjuvant activity of bacterial peptidoglycan derivatives*. Biochem Biophys Res Commun. (1974) 59: 1317-1325.

- Englund G, Theodore TS, Freed EO, Engleman A and Martin MA, *Integration is required for productive infection of monocyte-derived macrophages by human immunodeficiency virus type 1*. J Virol. (1995) 69: 3216-3219.

Fortin JF, Cantin R and Tremblay MJ, *T cells expressing activated LFA-1 are more susceptible to infection with human immunodeficiency virus type 1 particles bearing host-encoded ICAM*. J Virol. (1998) 72: 2105-2112.

- Gao F, Yue L, Craig S, Thornton CL, Robertson DL, McCutchan FE, Bradac JA, Sharp PM and Hahn BH, *Genetic variation of HIV type 1 in four World Health Organization-sponsored vaccine evaluation sites: generation of functional envelope (glycoprotein 160) clones representative of sequence subtypes A, B, C and E*. WHO Network for HIV Isolation and Characterization. AIDS Res Hum Retroviruses (1994)
 25 10: 1359-1368.

Ghiara JB, Stura EA, Stanfield RL, Profy AT and Wilson IA, *Crystal structure of the principal neutralization site of HIV*. Science (1994) 264: 82-85.

Higgins DG and Sharp PM, *Fast and sensitive multiple sequence alignments on a microcomputer*. CABIOS (1989) 5: 151-153.

Hioe CE, Xu S, Chigurupati P, Burda S, Williams C, Gorny MK and Zolla-Pazner S, *Neutralization of HIV-1 primary isolates by polyclonal and monoclonal human antibodies*. *Int Immunol.* (1997) 9: 1281-1290.

Korbert BT, MacInnes K, Smith RF and Myers G, *Mutational trends in V3 loop protein sequences observed in different genetic lineages of human immunodeficiency virus type 1*. *J Virol.* (1994) 68: 6730-6744.

Landau NR and Littman DR, *Packaging system for rapid production of murine leukemia virus vectors with variable tropism*. *J Virol.* (1992) 66: 5110-5113.

Lennette EH. "General principles underlying laboratory diagnosis of viral and rickettsial infections" in: Lennette EH and Schmidt MJ, *Diagnostic Procedures of Viral and Rickettsial Disease* (New York, American Public Health Association, 1964) pp. 45-

Liou HC, Sha WC, Scott ML and Baltimore D, *Sequential induction of NF-kappa B/Rel family proteins during B-cell terminal differentiation*. *Mol Cell Biol.* (1994) 14: 5349-5359.

Luciw PA. in: Fields BN, Knipe DM and Howley PM, *Fields Virology*, 3d ed, (Philadelphia, Lippincott-Raven, 1996), pp 1881-952.

Lusso P, Cocchi F, Balotta C, Markham PD, Louie A, Farci P, Pal R, Gallo RC and Reitz MS, *Growth of macrophage-tropic and primary human immunodeficiency virus type 1 (HIV-1) isolates in a unique CD4+ T-cell clone (PM1): failure to downregulate CD4 and to interfere with cell-line HIV-1*. *J Virol.* (1995) 69: 3712-3720.

Mackett M, Smith GL and Moss B, *General method for production and selection of infectious vaccinia virus recombinants expressing foreign genes*. *J Virol.* (1984) 49: 857-864.

Mann DL, O'Brien SJ, Gilbert DA, Reid Y, Popovic M, Read-Connole E, Gallo RC and Gazdar AF, *Origin of the HIV-susceptible human CD4+ cell line H9*. *AIDS Res Hum Retroviruses* (1989) 5: 253-255.

Mascola J, Louwagie J, McCutchan FE, Fischer CL, Hegerich PA, Wagner KF, Fowler AK, McNeil JG and Burke DS, *Two antigenically distinct subtypes of Human Immunodeficiency Virus Type 1: Viral genotype predicts neutralization serotype*. *J Infect Dis.* (1994) 169: 48-54.

- Mather JP, *Establishment and characterization of two distinct mouse testicular epithelial cell lines*. Biol Reprod. (1980) 23: 243-252.
-
- Mather JP, Zhuang LZ, Perez-Infante V and Phillips DM, *Culture of testicular cells in hormone-supplemented serum-free medium*. Ann NY Acad Sci. (1982) 383: 44-68.
- 5 Merrifield RB, *Automated synthesis of peptides*. Science (1965) 150: 178-185.
- Moore JP, Sattentau QJ, Yoshiyama H, Thali M, Charles M, Sullivan N, Poon S-W, Fung MS, Traincard F, Pinkus M, Robey G, Robinson JE, Ho DD and Sodroski J, *Probing the structure of the V2 domain of human immunodeficiency virus type 1 surface glycoprotein gp120 with a panel of eight monoclonal antibodies: human immune response to the V1 and V2 domains*. J Virol. (1993) 67: 6136-6151.
- 10 Montefiori DC, Pantaleo G, Fink LM, Zhou JT, Zhou JY, Bilska M, Miralles GD and Fauci AS, *Neutralizing and infection-enhancing antibody responses to human immunodeficiency virus type 1 in long-term nonprogressors*. J Infect Dis. (1996) 60-67.
- 15 Moore JP, Cao Y, Leu J, Qin L, Korber B and Ho DD. *Inter- and intracade neutralization of human immunodeficiency virus type 1: genetic clades do not correspond to neutralization serotypes but partially correspond to gp120 antigenic serotypes*. J Virol. (1996) 70: 427-44.
- Muster T, Stein F, Purtscher M, Trkola A, Klima A, Himmler G, Ruker F and Katinger H, *A conserved neutralization epitope on gp41 of human immunodeficiency virus type 1*. J Virol. (1993) 67: 6642-6647.
- 20 Myers EW and Miller W, *Optimal alignments in linear space*. CABIOS (1988) 4: 11-17.
- Myers G, Berzofsky JA, Korber B, Smith RF and Pavlakis GN, *Human retroviruses and AIDS 1992*. Los Alamos National Laboratory, Los Alamos, New Mexico, 1993.
- 25 Overbaugh J and Rudensey LM. *Alterations in potential sites for glycosylation predominate during evolution of the simian immunodeficiency virus envelope gene in macaques*. J Virol. (1992) 66: 5937-5948.

Park EJ, Vujcic LJ, Anand R, Theodore TS and Quinnan GV, *Mutations in both gp120 and gp41 are responsible for the broad neutralization resistance of variant HIV-1* *mn to antibodies directed at v3 and non-v3 epitopes*, J Virol. (1998) 72: 7099-7107.

Plantier JC, Le Pogam S, Poisson F, Buzelay L, Lejeune B, Barin F, *Extent of*
5 *antigenic diversity in the V3 region of the surface glycoprotein, gp120, of human*
immunodeficiency virus type 1 group M and consequences for serotyping. J Virol. (1998)
72: 677-83.

Quinnan G, Zhang P, Fu D, Dong M and Margolick J, *Evolution of neutralizing*
antibody response against hiv-1 virions and pseudovirions in multicenter aids cohort
10 *study participants.* AIDS Res Hum Retroviruses (1998) 14: 939-949.

Sabri F, Chiodi F and Fenyo EM, *Lack of correlation between V3 amino acid*
sequence and syncytium-inducing capacity of some HIV type 1 isolates. AIDS Res Hum
Retroviruses. (1996) 12: 855-858.

Saitou N and Nei M, *The neighbor-joining method: A new method for*
15 *reconstructing phylogenetic trees.* Mol. Biol. Evol (1987) 4: 406-425.

Schonning K, Jansson B, Olofsson S, Nielsen JO and Hansen JS. *Resistance to*
V3-directed neutralization caused by an N-linked oligosaccharide depends on the
quaternary structure of the HIV-1 envelope oligomer. Virol. (1996) 218: 134-140.

Seth A, Hodge DR, Thompson DM, Robinson L, Panayiotakis A, Watson DK and
20 Papas TS, *ETS family proteins activate transcription from HIV-1 long terminal repeat.*
AIDS Res Hum Retroviruses (1993) 9: 1017-1023.

Stuve LL, Brown-Shimer S, Pachl C, Naharian R, Diaz D and Burke RL,
Structure and expression of the herpes simplex virus type 2 glycoprotein gB gene. J
Virol. (1987) 61: 326-335.

Thali M, Charles M, Furman C, Cavacini L, Posner M, Robinson J and Sodroski
25 J, *Resistance to neutralization by broadly reactive antibodies to the human*
immunodeficiency virus type 1 gp120 glycoprotein conferred by a gp41 amino acid
change. J Virol. (1994) 68: 674-680.

Thali M, Furman C, Ho DD, Robinson J, Tilley S, Pinter A and Sodroski J,
30 *Discontinuous, conserved neutralization epitopes overlapping the CD4-binding region of*

human immunodeficiency virus type 1 gp120 envelope glycoprotein. J Virol. (1992) 66: 5635-5641.

Thali M, Moore JP, Furman C, Charles M, Ho DD, Robinson J and Sodroski J,

Characterization of conserved human immunodeficiency virus type 1 gp120

5 *neutralization epitopes exposed upon gp120-CD4.* J Virol. (1993) 67: 3978-3988.

Theodore TS, Englund G, Buckler-White A, Buckler CE, Martin MA and Peden KW, *Construction and characterization of a stable full-length macrophage-tropic HIV type 1 molecular clone that directs the production of high titers of progeny virions.* AIDS Res Hum Retroviruses (1996) 12: 191-194.

10 Trkola A, Purtscher M, Muster T, Ballaun C, Bauchacher A, Sullivan N, Srinivassan K, Sodroski J, Moore JP and Katinger H, *Human monoclonal antibody 2G12 defines a distinctive neutralization epitope of human immunodeficiency virus type 1.* J Virol. (1996) 70: 1100-1108.

15 Urlaub G and Chasin LA, *Isolation of Chinese hamster cell mutants deficient in dihydrofolate reductase activity.* Proc Natl Acad Sci USA (1980) 77: 4216-4220.

VanCott TC, Mascola JR, Kaminski RW, Kalyanaraman V, Hallberg PL, Burnett PR, Ulrich JT, Rechtman DJ and Birx DL, *Antibodies with specificity to native gp120 and neutralization activity against primary human immunodeficiency virus type 1 isolates elicited by immunization with oligomeric gp160.* J Virol. (1997) 71: 4319-4330.

20 Vujcic LK and Quinnan GV, *Preparation and characterization of human HIV type 1 neutralizing reference sera.* (1995) AIDS Res Hum Retroviruses. 11: 783-787.

Wrin T, Loh TP, Vennari JC, Schuitemaker H and Nunberg JH, *Adaptation to persistent growth in the H9 cell line renders a primary isolate of human immunodeficiency virus type 1 sensitive to neutralization by vaccine sera.* J Virol. (1995) 25 69: 39-48.

Zeng W, Regamey PO, Rose K, Wang Y and Bayer E. *Use of Fmoc-N-(2-hydroxy-4-methoxybenzyl)amino acids in peptide synthesis.* J Pept Res. (1997) 49: 273-279.

30 Zhang PF, Chen X, Fu DW, Margolick JB and Quinnan GV. *Primary virus envelope cross-reactivity of the broadening neutralizing antibody response during early chronic human immunodeficiency virus type 1 infection.* J Virol. (1999) 73: 5225-5230.

Zolla-Pazner S and Sharpe S. *A resting cell assay for improved detection of antibody-mediated neutralization of HIV type 1 primary isolates.* AIDS Res Hum

~~Retroviruses (1995) 11:1449-1458.~~

- Zwart G, Langedijk H, Van der Hoek L, de Jong JJ, Wolfs TF, Ramautarsing C,
5 Bakker M, De Ronde A and Goudsmit J, *Immunodominance and antigenic variation of the principal neutralization domain of HIV-1.* Virol. (1991) 181: 481-489.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50